arXiv:1604.00692v1 [cond-mat.dis-nn] 3 Apr 2016

A Lloyd-model generalization: Conductance fluctuations in one-dimensional disordered systems

J. A. Méndez-Bermúdez,[1] A. J. Martínez-Mendoza,[1,2] V. A. Gopar,[3] and I. Varga[2]
[1] *Instituto de Física, Benemérita Universidad Autónoma de Puebla, Apartado Postal J-48, Puebla 72570, Mexico*
[2] *Elméleti Fizika Tanszék, Fizikai Intézet, Budapesti Műszaki és Gazdaságtudományi Egyetem, H-1521 Budapest, Hungary*
[3] *Departamento de Física Teórica, Facultad de Ciencias, and BIFI, Universidad de Zaragoza, Pedro Cerbuna 12, E-50009, Zaragoza, Spain*

(Dated: April 5, 2016)

We perform a detailed numerical study of the conductance G through one-dimensional (1D) tight-binding wires with on-site disorder. The random configurations of the on-site energies ϵ of the tight-binding Hamiltonian are characterized by long-tailed distributions: For large ϵ, $P(\epsilon) \sim 1/\epsilon^{1+\alpha}$ with $\alpha \in (0, 2)$. Our model serves as a generalization of 1D Lloyd's model, which corresponds to $\alpha = 1$. First, we verify that the ensemble average $\langle -\ln G \rangle$ is proportional to the length of the wire L for all values of α, providing the localization length ξ from $\langle -\ln G \rangle = 2L/\xi$. Then, we show that the probability distribution function $P(G)$ is fully determined by the exponent α and $\langle -\ln G \rangle$. In contrast to 1D wires with standard white-noise disorder, our wire model exhibits bimodal distributions of the conductance with peaks at $G = 0$ and 1. In addition, we show that $P(\ln G)$ is proportional to G^β, for $G \to 0$, with $\beta \leq \alpha/2$, in agreement to previous studies.

PACS numbers: 72.10.-d, 72.15.Rn, 73.21.Hb

I. INTRODUCTION AND MODEL

The recent experimental realizations of the so-called Lévy glasses [1] as well as "Lévy waveguides" [2] has refreshed the interest in the study of systems characterized by Lévy-type disorder (see for example Refs. [3–15]). That is, disorder characterized by random variables $\{\epsilon\}$ whose density distribution function exhibits a slow decaying tail:

$$P(\epsilon) \sim \frac{1}{\epsilon^{1+\alpha}} \ , \tag{1}$$

for large x, with $0 < \alpha < 2$ (this kind of probability distributions are known as α-stable distributions [16]). In fact, the study of this class of disordered systems dates back to Lloyd [17], who studied spectral properties of a three-dimensional (3D) lattice described by a 3D tight-binding Hamiltonian with Cauchy-distributed on-site potentials [which corresponds to the particular value $\alpha = 1$ in Eq. (1)]. Since then, a considerable number of works have been devoted to the study of spectral, eigenfunction, and transport properties of Lloyd's model in its original 3D setup [18–27] and in lower dimensional versions [26–43].

Of particular interest is the comparison between the one-dimensional (1D) Anderson model (1DAM) [44] and the 1D Lloyd's model, since the former represents the most prominent model of disordered wires [45]. Indeed, both models are described by the 1D tight-binding Hamiltonian:

$$H = \sum_{n=1}^{L} [\epsilon_n \mid n\rangle \langle n \mid - \nu_{n,n+1} \mid n\rangle \langle n+1 \mid - \nu_{n,n-1} \mid n\rangle \langle n-1 \mid] \ ; \tag{2}$$

where L is the length of the wire given as the total number of sites n, ϵ_n are random on-site potentials, and $\nu_{n,m}$ are the hopping integrals between nearest neighbors (which are set to a constant value $\nu_{n,n\pm1} = \nu$). However, while for the standard 1DAM (with white-noise on-site disorder $\langle \epsilon_n \epsilon_m \rangle = \sigma^2 \delta_{nm}$ and $\langle \epsilon_n \rangle = 0$) the on-site potentials are characterized by a finite variance $\sigma^2 = \langle \epsilon_n^2 \rangle$ (in most cases the corresponding probability distribution function $P(\epsilon)$ is chosen as a box or a Gaussian distribution), in the Lloyd's model the variance σ^2 of the random on-site energies ϵ_n diverges since they follow a Cauchy distribution.

It is also known that the eigenstates Ψ of the *infinite* 1DAM are exponentially localized around a site position n_0 [45]:

$$|\Psi_n| \sim \exp\left(-\frac{|n - n_0|}{\xi}\right) \ ; \tag{3}$$

where ξ is the eigenfunction localization length. Moreover, for weak disorder ($\sigma^2 \ll 1$), the only relevant parameter for describing the statistical properties of the transmission of the *finite* 1DAM is the ratio L/ξ [46], a fact known as single parameter scaling. The above exponential localization of eigenfunctions makes the transmission or dimensionless conductance G exponentially small, i.e., [47]

$$\langle -\ln G \rangle = \frac{2L}{\xi} \ ; \tag{4}$$

thus, this relation can be used to obtain the localization length. Remarkably, it has been shown that Eq. (4) is also valid for the 1D Lloyd's model [41] implying a single parameter scaling, see also [38].

It is also relevant to mention that studies of transport quantities through 1D wires with Lévy-type disorder, different from the 1D Lloyd's model, have been reported. For example, wires with scatterers randomly

spaced along the wire according to a Lévy-type distribution were studied in Refs. [3, 4, 48, 49]. Concerning the conductance of such wires, a prominent result reads that the corresponding probability distribution function $P(G)$ is fully determined by the exponent α of the power-law decay of the Lévy-type distribution and the average (over disorder realizations) $\langle -\ln G\rangle$ [48, 49]; i.e., all other details of the disorder configuration are irrelevant. In this sense, $P(G)$ shows *universality*. Moreover, this fact was already verified experimentally in microwave random waveguides [2] and tested numerically using the tight-binding model of Eq. (2) with $\epsilon_n = 0$ and off-diagonal Lévy-type disorder [50] (i.e., with $\nu_{n,m}$ in Eq. (2) distributed according to a Lévy-type distribution).

It is important to point out that 1D tight-binding wires with power-law distributed random on-site potentials, characterized by power-laws different from $\alpha = 1$ (which corresponds to the 1D Lloyd's model), have been scarcely studied; for a prominent exception see [41]. Thus, in this paper we undertake this task and study numerically the conductance though disordered wires defined as a generalization of the 1D Lloyd's model as follows. We shall study 1D wires described by the Hamiltonian of Eq. (2) having constant hopping integrals, $\nu_{n,n\pm1} = \nu = 1$, and random on-site potentials ϵ_n which follow a Lévy-type distribution with a long tail, like in Eq. (1) with $0 < \alpha < 2$. We name this setup the 1DAM with Lévy-type on-site disorder. We note that when $\alpha = 1$ we recover the 1D Lloyd's model.

Therefore, in the following section we shall show that (i) the conductance distribution $P(G)$ is fully determined by the power-law exponent α and the ensemble average $\langle -\ln G\rangle$; (ii) for $\alpha \leq 1$ and $\langle -\ln G\rangle \sim 1$, bimodal distributions for $P(G)$ with peaks at $G \sim 0$ and $G \sim 1$ are obtained, revealing the coexistence of insulating and ballistic regimes; and (iii) the probability distribution $P(\ln G)$ is proportional to G^β, for vanishing G, with $\beta \leq \alpha/2$.

II. RESULTS AND DISCUSSION

Since we are interested in the conductance statistics of the 1DAM with Lévy-type on-site disorder we have to define first the scattering setup we shall use: We open the isolated samples described above by attaching two semi-infinite single channel leads to the border sites at opposite sides of the 1D wires. Each lead is also described by a 1D semi-infinite tight-binding Hamiltonian. Using the Heidelberg approach [51] we can write the transmission amplitude through the disordered wires as $t = -2i\sin(k)\,\mathcal{W}^T(E - \mathcal{H}_{\rm eff})^{-1}\mathcal{W}$, where $k = \arccos(E/2)$ is the wave vector supported in the leads and $\mathcal{H}_{\rm eff}$ is an effective non-hermitian Hamiltonian given by $\mathcal{H}_{\rm eff} = H - e^{ik}\mathcal{W}\mathcal{W}^T$. Here, $\mathcal{W}$ is a $L \times 1$ vector that specifies the positions of the attached leads to the wire. In our setup, all elements of $\mathcal{W}$ are equal to zero except $\mathcal{W}_{11}$ and $\mathcal{W}_{L1}$ which we set to unity (i.e., the leads are attached to the wire with a strength equal to the inter-site hopping amplitudes: $\nu = 1$). Also, we have fixed the energy at $E = 0$ in all our calculations, although the same conclusions are obtained for $E \neq 0$. Then, within a scattering approach to the electronic transport, we compute the dimensionless conductance as [52] $G = |t|^2$.



FIG. 1: (a) Average logarithm of the conductance $\langle -\ln G\rangle$ as a function of L for the 1DAM with Lévy-type on-site disorder (symbols). Dashed lines are the fittings of the data with Eq. (4) used to extract ξ. (b) $\langle\langle(-\ln G)^2\rangle\rangle$ as a function of L (symbols). Dashed lines are fittings of the data with the function $\langle\langle(-\ln G)^2\rangle\rangle = 4c_2 L$, see Eq. (5). In both panels $\alpha = 1/10$, $1/5$, $1/2$, 1, and $3/2$ (from top to bottom). Each point was calculated using 10^4 disorder realizations. $E = 0$ was used.

First, we present in Fig. 1(a) the ensemble average $\langle -\ln G\rangle$ as a function of L for the 1DAM with Lévy-type disorder for several values of α. It is clear from this figure that $\langle -\ln G\rangle \propto L$ for all the values of α we consider here. Therefore, we can extract the localization length ξ by fitting the curves $\langle -\ln G\rangle$ vs. L with Eq. (4); see dashed lines in Fig. 1(a). This behavior should be contrasted to the case of 1D wires with off-diagonal Lévy-type disorder [53] which shows the dependence $\langle -\ln G\rangle \propto L^{1/2}$ when $\alpha = 1/2$ at $E = 0$ [50].

Also, we have confirmed that the cumulants $\langle\langle(-\ln G)^k\rangle\rangle$ obey a linear relation with the wire length [41, 54], i.e.,

$$\lim_{L\to\infty} \frac{\langle\langle(-\ln G)^k\rangle\rangle}{L} = 2^k c_k \ , \qquad (5)$$

where the coefficients c_k, with $c_1 \equiv \xi^{-1}$, characterize the Lyapunov exponent of a generic 1D tight-binding wire with on-site disorder. We have verified the above relation, Eq. (5), for $k = 1$, 2, and 3; as an example in Fig. 1(b) we present the results for $\langle\langle(-\ln G)^2\rangle\rangle$ as a



FIG. 2: (Color online) Conductance distribution $P(G)$ for the 1DAM with Lévy-type disorder (histograms). Each panel correspond to a fixed value of $\langle -\ln G\rangle$: (a) $\langle -\ln G\rangle = 20$, (b) $\langle -\ln G\rangle = 2$, (c) $\langle -\ln G\rangle = 1$, (d) $\langle -\ln G\rangle = 2/3$, (e) $\langle -\ln G\rangle = 1/2$, and (f) $\langle -\ln G\rangle = 1/5$. In each panel we include histograms for several values of α, where α increases in the arrow direction. $E = 0$ was used. Each histogram was calculated using 10^6 disorder realizations. The red dashed lines are the theoretical predictions of $P(G)$ for the 1DAM with white noise disorder $P_{\rm WN}(G)$ corresponding to the particular value of $\langle -\ln G\rangle$ of each panel.



FIG. 3: (Color online) Conductance distribution $P(G)$ for the 1DAM with Lévy-type on-site disorder. Each panel corresponds to a fixed value of $\langle -\ln G\rangle$: (a) $\langle -\ln G\rangle = 1$, (b) $\langle -\ln G\rangle = 3/4$, (c) $\langle -\ln G\rangle = 1/2$, and (d) $\langle -\ln G\rangle = 1/4$. In each panel we include histograms for $\alpha = 1/4$, $1/2$, $3/4$, and 1; where α increases in the arrow direction. $E = 0$ was used. For each value of α we present two histograms using different Lévy-type density distributions of on-site disorder: $\rho_1(\epsilon)$ in red and $\rho_2(\epsilon)$ in black; see Eqs. (8) and (9) in [57]. Each histogram was calculated using 10^6 disorder realizations.

function of L for different values of α. The dashed lines are fittings of the numerical data (open dots) with the function $\left\langle\left\langle(-\ln G)^2\right\rangle\right\rangle = 4c_2 L$, see Eq. (5), which can be used to extract the higher order coefficient c_2.

Now, in Fig. 2 we show different conductance distributions $P(G)$ for the 1DAM with Lévy-type on-site disorder for fixed values of $\langle -\ln G\rangle$; note that fixed $\langle -\ln G\rangle$ means fixed ratio L/ξ. Several values of α are reported in each panel. We can observe that for fixed $\langle -\ln G\rangle$, by increasing α the conductance distribution evolves towards the $P(G)$ corresponding to the 1DAM with white noise disorder, $P_{\rm WN}(G)$, as expected. The curves for $P_{\rm WN}(G)$ are included as a reference in all panels of Fig. 2 as red dashed lines [55]. In fact, $P(G)$ already corresponds to $P_{\rm WN}(G)$ once $\alpha = 2$.

We recall that for 1D tight-binding wires with off-diagonal Lévy-type disorder $P(G)$ is fully determined by the exponent α and the average $\langle -\ln G\rangle$ [50]. It is therefore pertinent to ask whether this property also holds for *diagonal* Lévy-type disorder. Thus, in Fig. 3 we show $P(G)$ for the 1DAM with Lévy-type on-site disorder for several values of α, where each panel corresponds to a fixed value of $\langle -\ln G\rangle$. For each combination of $\langle -\ln G\rangle$ and α we present two histograms (in red and black) corresponding to wires with on-site random potentials $\{\epsilon_n\}$ characterized by two *different* density distributions [57], but with the same exponent α of their corresponding power-law tails. We can see from Fig. 3 that for each value of α the histograms (in red and black) fall on the top of each other, which is an evidence that the conductance distribution $P(G)$ for the 1DAM with Lévy-type on-site disorder is invariant once α and $\langle -\ln G\rangle$ are fixed; i.e., $P(G)$ displays a universal statistics.

Moreover, we want to emphasize the coexistence of insulating and ballistic regimes characterized, respectively, by the two prominent peaks of $P(G)$ at $G = 0$ and $G = 1$. This behavior, which is more evident for $\langle -\ln G\rangle \sim 1$ and $\alpha \leq 1$ (see Figs. 2 and 3), is not observed in 1D wires with white-noise disorder (see for example the red dashed curves in Fig. 2). This coexistence of opposite transport regimes has been already reported in systems with anomalously localized states: 1D wires with obstacles randomly spaced according to Lévy-type density distribution [48, 50] as well as in the so-called random-mass Dirac model [58].

Finally, we study the behavior of the tail of the distribution $P(\ln G)$. Thus, using the same data of Fig. 3,



FIG. 4: (Color online) Probability distribution functions $P(\ln G)$ for the 1DAM with Lévy-type on-site disorder. Same parameters as in Fig. 3. We just recall that in each panel we included histograms for $\alpha = 1/4$, $1/2$, $3/4$, and 1. Here, α increases in the arrow direction.



FIG. 5: The exponent β, see Eq. (6), as a function of α for $\langle -\ln G\rangle = 1/10$ (circles), 1 (diamonds), and 10 (triangles). The dashed line corresponds to $\beta = \alpha/2$. β was obtained from power-law fittings of the tails of the histograms of $P(\ln G)$ in the interval $P(\ln G) \in [10^{-5}, 10^{-3}]$.

in Fig. 4 we plot $P(\ln G)$. As expected, since $P(G)$ is determined by α and $\langle -\ln G\rangle$, we can see that $P(\ln G)$ is invariant once those two quantities (α and $\langle -\ln G\rangle$) are fixed (red and black histograms fall on top of each other). Moreover, from Fig. 4 we can deduce a power-law behavior:

$$P(\ln G) \propto G^{\beta} \tag{6}$$

for $G \to 0$ when $\alpha < 2$. For $\alpha = 2$, $P(\ln G)$ displays a log-normal tail (not shown here), expected for 1D systems in the presence of Anderson localization. Actually, the behavior (6) was already anticipated in [41] as $P(G) \sim G^{-(2-\lambda)/2}$ for $G \to 0$ with $\lambda < \alpha$; which in our study translates as $P(\ln G) \propto G^{\lambda/2}$ (since $P(\ln G) = GP(G)$) with $\lambda/2 \equiv \beta \leq \alpha/2$. Indeed, we have validated the last inequality in Fig. 5 where we report the exponent β obtained from power-law fittings of the tails of the histograms of $P(\ln G)$. In addition, we have observed that the value of β depends on the particular value of $\langle -\ln G\rangle$ characterizing the corresponding histogram of $P(\ln G)$. Also, from Fig. 5 we note that $\beta \approx \alpha/2$ as the value of $\langle -\ln G\rangle$ decreases.

III. CONCLUSIONS

In this work we have studied the conductance G through a generalization of Lloyd's model in one dimension: We consider one-dimensional (1D) tight-binding wires with on-site disorder following a Lévy-type distribution, see Eq. (1), characterized by the exponent α of the power-law decay. We have verified that different cumulants of the variable $\ln G$ decrease linearly with the length wire L. In particular, we were able to extract the eigenfunction localization length ξ from $\langle -\ln G\rangle = 2L/\xi$. Then, we have shown some evidence that the probability distribution function $P(G)$ is invariant, i.e., fully determined, once α and $\langle -\ln G\rangle$ are fixed; in agreement with other Lévy-disordered wire models [2, 48–50]. We have also reported the coexistence of insulating and ballistic regimes, evidenced by peaks in $P(G)$ at $G = 0$ and $G = 1$; these peaks are most prominent and commensurate for $\langle -\ln G\rangle \sim 1$ and $\alpha \leq 1$. Additionally we have shown that $P(\ln G)$ develops power-law tails for $G \to 0$, characterized by the power-law β (also invariant for fixed α and $\langle -\ln G\rangle$) which, in turn, is bounded from above by $\alpha/2$. This upper bound of β implies that the smaller the value of α the larger the probability to find vanishing conductance values in our Lévy-disordered wires.

Acknowledgments

J.A.M.-B. and A.J.M.-M. thank F. M. Izrailev and N. M. Makarov for useful comments. J.A.M.-B. and A.J.M.-M. also thank FAPESP (Grant No. 2014/25997-0), CONACyT (Grants No. I0010-2014-246246 and No. CB-2013-220624), VIEP-BUAP (Grant No. MEBJ-EXC15-I), and PIFCA (Grant No. BUAP-CA-169) for financial support. V.A.G. acknowledges support from MINECO (Spain) under the Project number FIS2012-35719-C02-02.

[1] P. Barthelemy, J. Bertolotti, and D. S. Wiersma, Nature **453**, 495 (2008).

[2] A. A. Fernandez-Marin, J. A. Mendez-Bermudez, J. Carbonell, F. Cervera, J. Sanchez-Dehesa, and V. A. Gopar, Phys. Rev. Lett. **113**, 233901 (2014).

[3] C. W. J. Beenakker, C. W. Groth, and A. R. Akhmerov, Phys. Rev. B **79**, 024204 (2009).

[4] R. Burioni, L. Caniparoli, and A. Vezzani, Phys. Rev. E **81**, 060101(R) (2010).

[5] A. Eisfeld, S. M. Vlaming, V. A. Malyshev, and J. Knoester, Phys. Rev. Lett. **105**, 137402 (2010).

[6] J. Bertolotti, K. Vynck, L. Pattelli, P. Barthelemy, S. Lepri, D. S. Wiersma, Adv. Functional Materials **20**, 965 (2010).

[7] P. Barthelemy, J. Bertolotti, K. Vynck, S. Lepri, D. S. Wiersma, Phys. Rev. E **82**, 011101 (2010).

[8] M. Burresi, V. Radhalakshmi, R. Savo, J. Bertolotti, K. Vynck, and D. S. Wiersma, Phys. Rev. Lett. **108**, 110604 (2012).

[9] C. W. Groth, A. R. Akhmerov, and C. W. J. Beenakker, Phys. Rev. E **85**, 021138 (2012).

[10] R. Burioni, S. di Santo, S. Lepri, and A. Vezzani, Phys. Rev. E **86**, 031125 (2012).

[11] S. M. Vlaming, V. A. Malyshev, A. Eisfeld, and J. Knoester, J. Chem. Phys. **138**, 214316 (2013).

[12] R. Burioni, E. Ubaldi, and A. Vezzani, Phys. Rev. E **89**, 022135 (2014).

[13] P. Bernabo, R. Burioni, S. Lepri, and A. Vezzani, Chaos Solitons Fractals **67**, 11 (2014).

[14] S. S. Zakeri, S. Lepri, and D. S. Wiersma, Phys. Rev. E **91**, 032112 (2015).

[15] A. G. Ardakani and M. G. Nezhadhaghighi, J. Opt. **17**, 105601 (2015).

[16] V. V. Uchaikin and V. M. Zolotarev, *Chance and Stability. Stable Distributions and their Applications* (VSP, Utrecht, 1999).

[17] P. Lloyd, J. Phys. C **2**, 1717 (1969).

[18] M. Saitoh, Phys. Lett. A **33**, 44 (1970); Progr. Theor. Phys. **45**, 746 (1971).

[19] A. P. Kumar and G. Baskaran, J. Phys. C **6**, L399 (1973).

[20] K. Hoshino, Phys. Lett. A **56**, 133 (1976).

[21] W. R. Bandy and A. J. Glick, Phys. Rev. B **16**, 2346 (1977).

[22] S. Kivelson and C. D. Gelatt, Phys. Rev. B **20**, 4167 (1979).

[23] B. Simon, Phys. Rev. B **27**, 3859 (1983).

[24] D. E. Rodrigues and J. F. Weisz, Phys. Rev. B **34**, 2306 (1986).

[25] E. Kolley and W. Kolley, J. Phys. C **21**, 6099 (1988).

[26] R. Johnston and H. Kunz, J. Phys. C **16**, 4565 (1983).

[27] D. E. Rodrigues, H. M. Pastawski, and J. F. Weisz, Phys. Rev. B **34**, 8545 (1986).

[28] D. J. Thouless, J. Phys. C **5**, 77 (1972).

[29] K. Ishii, Suppl. Progr. Theor. Phys. **53**, 77 (1973).

[30] R. Abou-Chacra and D. J. Thouless, J. Phys. C **7**, 65 (1974).

[31] D. J. Thouless, J. Phys. C **16**, L929 (1983).

[32] A. MacKinnon, J. Phys. C **17**, L289 (1984).

[33] M. O. Robbins and B. Koiller, Phys. Rev. B **32**, 4576 (1985).

[34] D. L. Shepelyansky, Phys. Rev. Lett. **56**, 677 (1986).

[35] S. Fishman, R. E. Prange, and M. Griniasty, Phys. Rev. A **39**, 1628 (1989).

[36] G. Casati, I. Guarneri, F. Izrailev, S. Fischman, and L. Molinari, J. Phys.: Condens. Matter **4**, 149 (1992).

[37] C. Mudry, P. W. Brouwer, B. I. Halperin, V. Gurarie, and A. Zee, Phys. Rev. B **58**, 13539 (1998).

[38] L. I. Deych, A. A. Lisyansky, and B. L. Altshuler, Phys. Rev. Lett. **84**, 2678 (2000); Phys. Rev. B **64**, 224202 (2001).

[39] D. M. Gangardt and S. Fishman, Phys. Rev. B **63**, 045106 (2001).

[40] C. Fuchs and R. v. Baltz, Phys. Rev. B **63**, 085318 (2001).

[41] M. Titov and H. Schomerus, Phys. Rev. Lett. **91**, 176601 (2003).

[42] D. Roy and N. Kumar, Phys. Rev. B **76**, 092202 (2007).

[43] G. G. Kozlov, Theor. Math. Phys. **171**, 531 (2012).

[44] P. W. Anderson, Phys. Rev. **109**, 1492 (1958).

[45] *50 Years of Anderson Localization*, E. Abrahams, ed. (World Scientific, Singapore, 2010).

[46] P. W. Anderson, D. J. Thouless, E. Abrahams, and D. S. Fisher, Phys. Rev. B **22**, 3519 (1980).

[47] I. M. Lifshits, S. A. Gredeskul, and L. A. Pastur, *Introduction to the theory of disordered systems* (Willey, New York, 1988).

[48] F. Falceto and V. A. Gopar, Europhys. Lett. **92**, 57014 (2010).

[49] A. A. Fernandez-Marin, J. A. Mendez-Bermudez, and V. A. Gopar, Phys. Rev. A **85**, 035803 (2012).

[50] I. Amanatidis, I. Kleftogiannis, F. Falceto, and V. A. Gopar, Phys. Rev. B **85**, 235450 (2012).

[51] C. Mahaux and H. A Weidenmüller, *Shell Model Approach in Nuclear Reactions*, (North-Holland, Amsterdam,1969); J. J. M. Verbaarschot, H. A. Weidenmüller, and M. R. Zirnbauer, Phys. Rep. **129**, 367 (1985); I. Rotter, Rep. Prog. Phys. **54**, 635 (1991).

[52] R. Landauer, IBM J. Res. Dev. **1**, 223 (1957); **32**, 336 (1988); M. Buttiker, Phys. Rev. Lett. **57**, 1761 (1986); IBM J. Res. Dev. **32**, 317 (1988).

[53] It is pertinent to remark that the dependence $\langle -\ln G\rangle \propto L^{1/2}$, when $\alpha = 1/2$ at $E = 0$, reported in [50] for 1D wires with off-diagonal Lévy-type disorder was observed when the wire length L was defined as the total sum of the hopping integrals $L = \sum_n \nu_{n,n+1}$.

[54] H. Schomerus and M. Titov, Eur. Phys. J. B **35**, 421 (2003); Phys. Rev. B **67**, 100201(R) (2003).

[55] Using the results in Ref. [56], Eq. (2) of that work, $P_{\rm WN}(G)$ is given by

$$P_{\rm WN}(G) = C\sqrt{\frac{\operatorname{acosh}\left(1/\sqrt{G}\right)}{G^3\sqrt{1-G}}}\exp\left[-\frac{1}{s}\operatorname{acosh}^2\left(\frac{1}{\sqrt{G}}\right)\right], \tag{7}$$

where C is a normalization constant and $s = L/\ell$, ℓ being the mean free path. The parameter s can the obtained numerically from the ensemble average $\langle \ln G\rangle = -L/\ell$.

[56] I. Kleftogiannis, I. Amanatidis, V. A. Gopar, Phys. Rev. B **88**, 205414 (2013).

[57] We have used the particular density distributions:

$$\rho_1(\epsilon) = \frac{1}{\Gamma(\alpha)}\left(\frac{1}{2}\right)^{\alpha}\frac{1}{\epsilon^{1+\alpha}}\exp\left(-\frac{1}{2\epsilon}\right) \tag{8}$$

and

$$\rho_2(\epsilon) = \frac{\alpha}{(1+\epsilon)^{1+\alpha}} \ , \tag{9}$$

where Γ is the Euler gamma function.

[58] M. Steiner, Y. Chen, M. Fabrizio, and A. O. Gogolin, Phys. Rev. B **59**, 14848 (1999).